


Published: 2020-09-30

RADIOGRAPHIC FINDINGS OF A SYMPTOMATIC PNEUMOTHORAX DURING INTERCOSTAL NERVE INFILTRATION



Ivan Urits, MD¹
Ruben Schwartz, DO²
Alaa Abd-Elseyed, MD³
Omar Viswanath, MD⁴⁻⁶
Vwaire Orhurhu, MD¹
Alan D. Kaye, MD, PhD⁷
Cyrus Yazdi, MD¹

Fig. 1. (A) lateral, (B) magnified lateral, (C) anteroposterior, and (D) magnified anteroposterior radiographs demonstrating a 10% to 15% right pneumothorax. Important radiographic findings include a visible pleural edge and absent lung markings peripheral to this demarcation.

Intercostal nerve blocks are a relatively simple procedure that may be performed for the treatment of chronic pain conditions, such as postherpetic neuralgia and postsurgical incision pain. Although widely regarded as safe, intercostal nerve blocks among other thoracic procedures may lead to inadvertent pneumothorax. The

incidence of pneumothorax during intercostal nerve block can vary widely; historically, rates of 0.073% to 19% have been reported (1-3). More recently, in a study of 1,020 individual intercostal nerve blocks, pneumothorax occurred in 1.4% of procedures performed (4). Current standard for universal use of ultrasound guid-

From: ¹Department of Anesthesia, Critical Care, and Pain Medicine, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA; ²Department of Anesthesiology, Mount Sinai Medical Center of Florida, Miami, FL; ³Department of Anesthesiology, Chronic Pain Medicine, University of Wisconsin-Madison, Madison, WI; ⁴Valley Anesthesiology and Pain Consultants, Phoenix, AZ; ⁵Department of Anesthesiology, University of Arizona College of Medicine-Phoenix, Phoenix, AZ; ⁶Department of Anesthesiology, Creighton University School of Medicine, Omaha, NE; ⁷Department of Anesthesiology, Louisiana State University Health Sciences Center, New Orleans, LA

Corresponding Author: Ruben Schwartz, DO, E-mail: rubenschwartz@yahoo.com

Conflict of Interest: Each author certifies that he or she, or a member of his or her immediate family, has no commercial association (i.e., consultancies, stock ownership, equity interest, patent/licensing arrangements, etc.) that might pose a conflict of interest in connection with the submitted manuscript.

Disclaimer: There was no external funding in the preparation of this manuscript.

ance when performing an intercostal nerve block may further reduce the incidence of pneumothorax. Pneumothorax may have a delayed presentation, although often presents either during or immediately after the inciting procedure.

Chest radiography is the first diagnostic modality performed to assess pneumothorax as it is quick and inexpensive; of note, computed tomography offers greater sensitivity in assessing for a small pneumothorax. These radiographs (Fig. 1) demonstrate a 10% to 15% right-sided pneumothorax in a 55-year-old man who

underwent a right-sided ultrasound-guided intercostal nerve block at the T8 vertebral level. Immediately after the injection he reported shortness of breath and a change in oxygen saturation from 100% to 98%. As in the instance of this patient, observation alone is the treatment of choice for small pneumothorax with mild breathlessness (5). Markedly symptomatic patients should be admitted for further assessment and management. Despite this, providers should be ready to urgently diagnose and decompress a pneumothorax in the acutely presenting patient.

REFERENCES

1. Bartlett RW, Eastwood DW. Long-acting bilateral intercostal nerve block for upper abdominal surgery. *Surgery* 1952; 32:956-960.
2. McCleery RS, Zollinger R, Lenahan NE. A clinical study of the effect of intercostal nerve block with nupercaine in oil following upper abdominal surgery. *Surg Gynecol Obstet* 1948; 86:680-686.
3. Moore DC. Intercostal nerve block for postoperative somatic pain following surgery of thorax and upper abdomen. *Br J Anaesth* 1975; 47(suppl):284-286.
4. Shanti CM, Carlin AM, Tyburski JG. Incidence of pneumothorax from intercostal nerve block for analgesia in rib fractures. *J Trauma* 2001; 51:536-539.
5. MacDuff A, Arnold A, Harvey J; BTS Pleural Disease Guideline Group. Management of spontaneous pneumothorax: British Thoracic Society pleural disease guideline 2010. *Thorax* 2010; 65(suppl 2):ii18-31.